



J. Hoenicke
A. Nutz

30.4.2013
hand in until 7.5.2013, 10:15

Tutorials for Decision Procedures Exercise sheet 3

Exercise 1: Prenex Normal Form

Transform the following formula into prenex normal form:

$$\left(\forall z. \left((\forall x. q(x, z)) \rightarrow p(x, g(y), z) \right) \right) \wedge \neg(\forall z. \neg(\forall x. q(f(x, y), z)))$$

Exercise 2: Syntax and Semantics of FOL

- (a) Assume that a, b are constant symbols, f resp. g are function symbols of arity one resp. two, p is a predicate symbol of arity two, and x is a variable. For each of the following strings determine whether it is a *term*, an *atom*, a *literal*, or a *formula*. Note that it can be more than one of these, or none if it is syntactically incorrect.

- | | | |
|--------------------|------------------------------|-------------------------------------|
| (i) $f(a)$ | (v) $\neg p(a, b)$ | (ix) $p(a, b) \vee p(b, a)$ |
| (ii) $g(f(a), f)$ | (vi) $\exists a.p(a, b)$ | (x) $p \wedge \exists x.p(x, x)$ |
| (iii) $p(f(a), x)$ | (vii) $\exists x.p(x, f(a))$ | (xi) $\neg \exists x.p(a, b)$ |
| (iv) $g(x, f(x))$ | (viii) $p(x, p(x, x))$ | (xii) $\forall x.\exists x.p(x, x)$ |

- (b) For each of the following formulae give a satisfying interpretation. Assume that *equals* and *p* are binary predicates, *f* is a unary function, and *add* is a binary function.

- | | |
|----------------------------|--|
| (i) $equals(add(2, 2), 5)$ | (iii) $\exists y.\forall x.p(x, y)$ |
| (ii) $\forall x.p(x, x)$ | (iv) $\forall x.(p(x, f(x)) \wedge \neg p(f(x), x))$ |

Exercise 3: Semantic Tableaux for FOL

Use the semantic tableaux method to prove the validity of the following formulae.

- $(\forall x. (p(x) \rightarrow q(a))) \wedge (\exists x. p(x)) \rightarrow q(a)$
- $(\forall x. p(f(x))) \wedge (\forall y. (q(y) \rightarrow \neg p(f(y)))) \rightarrow \neg q(b)$
- $(\forall x, y. (p(x, y) \vee p(y, x))) \rightarrow \forall z.p(z, z)$
- $\forall y. \exists x. (p(x) \rightarrow p(y))$
- $\exists x. \forall y. (p(x) \rightarrow p(y))$