



J. Hoenicke  
A. Nutz

28.05.2013  
submit until 04.06.2013, 10:15

## Tutorials for Decision Procedures Exercise sheet 5

### Exercise 1: Integer Arithmetic

Consider the  $T_{\mathbb{Z}}$ -formula  $F : \exists x. \forall y. \neg(y + 1 = x)$ .

- Convert  $F$  into an equisatisfiable  $T_{\mathbb{N}}$ -formula  $G$ .
- Prove unsatisfiability of  $G$  using the semantic tableaux method. You may assume that associativity and commutativity of addition holds.
- Prove validity of the  $T_{\mathbb{N}}$ -formula  $\exists x. \forall y. \neg(y + 1 = x)$ .

### Exercise 2: Quantifier Elimination for $T_{\mathbb{Q}}$

Apply quantifier elimination to the following  $T_{\mathbb{Q}}$ -formulae:

- $\exists y. (x = 2y \wedge y < x)$
- $\forall y. (25 < x + 2y \vee x + 2y < 25)$
- $\forall x. \exists y. (y > x \wedge -y < x)$
- $\forall x. (x > 0 \iff \exists y. (x > y \wedge -x < y))$

### Exercise 3: Sufficient Set

For  $T_{\mathbb{Q}}$  the algorithm in the lecture examines terms  $\frac{s+t}{2}$  for all  $s, t \in S$ . Suppose we split up  $S$  in  $S_A, S_B, S_C$  depending on whether the term  $t$  comes from an (A)  $x < t$ , (B)  $t < x$ , or (C)  $x = t$  literal. Based on this distinction, give a smaller set of terms that still is sufficient.