



J. Hoenicke
A. Nutz

11.06.2012
submit until 18.06.2012, 10:15

Tutorials for Decision Procedures Exercise sheet 7

Exercise 1: Deciding $T_{\mathbb{Q}}$

Apply the Dutertre-de-Moura algorithm to decide the $T_{\mathbb{Q}}$ -satisfiability of the following $\Sigma_{\mathbb{Q}}$ -formulae: Give a satisfying $T_{\mathbb{Q}}$ -interpretation if it exists.

- (a) $x + 2y \geq 1 \wedge 2x + y \geq 1 \wedge x + y \leq \frac{1}{2}$
- (b) $x + 2y \geq 1 \wedge 2x + y \geq 1 \wedge x + y \leq 1$
- (c) $x + 2y > 1 \wedge 2x + y > 1 \wedge x + y < 1$
- (d) $x + 2y \geq 1 \wedge 2x + y \geq 1 \wedge x + y < \frac{2}{3}$

Exercise 2: Implementing the decision procedure for $T_{\mathbb{Q}}$ (8points)

Implement the Dutertre de Moura Algorithm from the lecture for input in the SMTLIB 2 format you already know.

Assume a conjunctive fragment where only atoms of the form $(\leq (+ t_1 \dots t_n) c_i)$ are asserted.

$t_i ::= (* c_i x_i) \mid (- x_i) \mid x_i$

$c_i ::= (- n) \mid n$

with n being a natural number and x_i being variable names.

On the lecture's website there is code which generates a tableau and the other necessary values from such input. There also is a readme-file with some technical instructions at the same place.